

PHYS 3038 Optics

L6 Geometrical Optics

Reading Material: Ch5.4-5.9



Shengwang Du




2015, the Year of Light



5.4.1


Planar Mirrors

(a)



(b)





The ability to rapidly redirect a beam of light by virtue of planar mirrors that has been utilized in the design of a single-lens reflex camera (see photo on page 11).




Figure 5.43 Inversions via reflection.





Figure 5.44 Rotation of a mirror and the concomitant angular displacement of a beam.



This tiny tiltable mirror (which is so small it can fit through the eye of a needle) is used to steer light beams in one of today's most important telecommunications devices. (Photo courtesy Lucent Technologies' Bell Laboratories.)

Aspherical Mirrors




Figure 5.45
A paraboloidal mirror.

$$OPL = \overline{W_1 A_1} + \overline{A_1 F} = \overline{W_2 A_2} + \overline{A_2 F}$$

$$\overline{W_1 A_1} + \overline{A_1 D_1} = \overline{W_2 A_2} + \overline{A_2 D_2}$$

Paraboloidal Mirror

co



Aspherical Mirrors





Figure 5.47 Hyperbolic and elliptical mirrors.



The 2.4-m-diameter hyperboloidal primary mirror of the Hubble Space Telescope. (Photo courtesy of NASA.)

5.4.3 Spherical Mirrors







Figure 5.49
Comparison of spherical and paraboloidal mirrors.

Mirror Formula




$$R < 0$$

Figure 5.50 A concave spherical mirror. Conjugate foci.


$$\frac{1}{s_o} + \frac{1}{s_i} = \frac{1}{f}$$

$$f = -\frac{R}{2}$$



Negative Focal Length

☞




$$R > 0$$

$$\frac{1}{s_o} + \frac{1}{s_i} = \frac{1}{f}$$

$$f = -\frac{R}{2}$$

Imagery with spherical mirrors

‘g’



$$\frac{1}{s_o} + \frac{1}{s_i} = \frac{1}{f}$$

$$f = -\frac{R}{2}$$

5.5.1 Dispersing Prisms






Figure 5.56 Geometry of a dispersing prism

$$\sin \theta_{i1} = n \sin \theta_{t1}$$

$$n \sin \theta_{i2} = \sin \theta_{t2}$$



Dispersing Prisms



$$n = \frac{\sin[(\delta_m + \alpha)/2]}{\sin \alpha/2}$$

Constant Deviation Dispersing Prisms




Figure 5.58 The Pellin–Broca prism.






Figure 5.59 The Abbe prism.

5.5.2 Reflecting Prisms

(internal reflection without dispersion)



Reflecting Prisms








Figure 5.63 The Dove prism.




3




Fiber Optics



$$NA = n_i \sin \theta_{max}$$



Colonoscope



Fiberoptical Communication



Charles K Kao
Nobel Prize in Physics in 2009

“for groundbreaking achievements concerning the transmission of light in fibers for optical communication”




Figure 5.72 The three major fiberoptic configurations and their index profiles. (a) Multimode step-index fiber. (b) Multimode graded-index fiber. (c) Single-mode step-index fiber.

5.7 Optical Systems




❖ Eyes

❖ Eyeglasses

❖ Microscope

❖ Camera

❖ Telescope









© 2004 MedicineNet, Inc.


Eye



Eyecllasses





Microscope



Camera



Telescope



Hubble Telescope

